

# 中国牙甲总科滨甲属昆虫志

蒲 蛰 龙

(中山大学生物系昆虫学教研组)

我国鞘翅目牙甲总科滨甲属(*Ochthebius*)的昆虫, 1939年以前的记录, 只有3种。1939年以后, 由我国各地陆续采集了不少这一属昆虫, 现将本属的昆虫进行分类学的研究, 使能进一步了解我国滨甲属昆虫的种类和分布。

截至目前止, 发现的滨甲属昆虫共有12种, 1亚种, 分隶6个亚属。

## 属的一般形态

体形长, 体色棕、棕黑或黑, 常有金属光泽。虫体背面常披有微小的短毛或平伏的长毛, 也有光滑无毛的。

头部一般有刻点, 刻点的粗细深浅因种类而异。点间光滑或粗糙。额头顶区有二显著的凹陷, 头的后部有一较小的凹陷。复眼突起, 甚显著。唇基颇大。口上缝一般稍弯。上唇大, 宽度约与唇基相等, 前缘完整, 或成深浅不等的弯缺或缺刻。触角短, 锤状, 9节, 基部4节光滑, 端部5节披毛, 成长形的锤状部。下颚鬚4节, 较触角短, 第1节短小, 第2节细长, 第3节长而较粗, 其端部尤粗大, 第4节短小, 在 *Henicocerus* 亚属尤甚。颜梯形或矩形, 通常有稀疏刻点, 前缘常有缺刻。

前胸背板可分为两部分, 中央部分称为盘状部, 两侧称为耳状部。盘状部与耳状部间由一条纵形或斜形的沟所分隔。前胸背板形状因种类和亚属的不同而异, 有横形的, 有近矩形的, 有近心形的。前胸背板的刻纹形状和数目也因种类和亚属的不同而异, 有成纵形沟, 横形沟, 斜形沟, 深的刻陷或浅的刻陷。有许多种类在盘状部有一中央纵沟, 两侧有前后刻陷各一对, 这些刻陷称为盘部前陷和盘部后陷。前胸背板前缘(包括盘状部及耳状部的前缘, 本文以后应用这个名词, 均包括这两部分)弯陷, 近于平直, 弯拱或略呈波状弯曲; 后缘一般略弯拱, 较前缘窄; 侧缘(包括盘状部和耳状部的侧缘, 本文叙述前胸背板侧缘, 均包括这两部分)自前部起, 或自中部起, 或自后部起缩窄, 缩窄部有透明膜围绕。

小盾片小, 三角形, 有刻纹或光滑。

前胸腹板短, 通常披有稀疏短毛, 在前足基节间的部分极窄。中胸腹板通常披有稀疏短毛, 由二纵形小隆脊纵分为三部分, 前部有一横形小隆脊, 在中足基节间的部分通常呈三角形。后胸腹板大, 有些种类全部光滑无毛, 有些全部披毛, 还有一些中部光滑, 其余部分披毛。

鞘翅长形或卵形, 隆拱, 有纵形条纹或由刻点所成的纵纹, 有时刻点混乱而不成条, 亦有无显著刻点的。翅基突缘通常明显。鞘翅后角通常宽圆, 缘折一般不达于鞘翅的末端, 但 *Henicocerus* 亚属的鞘翅缘折达于鞘翅末端。

足顏色較淡。前足及中足基节短柄状、錐状或筒状，粗糙，一般披柔毛，后足基节长，横置，粗糙，一般披柔毛。腿节光耀，披稀疏短毛或不披毛。脛节光耀，較长于或等长于腿节，通常有刺及刻点，端部較大。跗节 5 节，較脛节短，第 1 节極短，第 2 节也短，这两节互相紧接，分界極不明显，現为一节，第 3 节及第 4 节约等长，末节較长。爪纖細而弯。

腹部有 7 腹板，基部 5 腹板色較深而暗，披柔毛，第 6、第 7 腹板色淡而光耀，有时有稀疏的短毛或微小刻紋或刻点。

在中国發現的滨甲屬的亚屬共有 6 个，*Henicocerus*, *Aulacochthebius*, *Homalochthebius*, *Asiobates*, *Bothochius* 及 *Hymenodes*。

### 滨甲屬亚屬的检索表

- |   |                        |
|---|------------------------|
| 1(2) 鞘翅緣折达鞘翅的末端。下顎鬚末节很短.....                        | <i>Henicocerus</i>     |
| 2(1) 鞘翅緣折不达鞘翅的末端。下顎鬚末节較长.....                       |                        |
| 3(4) 前胸背板極寬而短，有二深橫沟。体形小，长约 1 毫米.....                | <i>Aulacochthebius</i> |
| 4(3) 前胸背板不大橫寬，一般無橫沟，有时有浅橫沟。体形較大，一般体长在 1.5 毫米以上..... |                        |
| 5(7) 前胸背板寬度显著地超过长度，側緣自后半部起向内縮窄.....                 |                        |
| 6(10) 前胸背板有一中央縱沟，盘状部一般無刻陷，如有刻陷，也不明显.....            | <i>Homalochthebius</i> |
| 7(5) 前胸背板呈心形，側緣自中部或自前半部起向内縮窄.....                   |                        |
| 8(9) 鞘翅的刻点不規則.....                                  | <i>Bothochius</i>      |
| 9(8) 鞘翅的刻点排成許多縱列紋.....                              | <i>Hymenodes</i>       |
| 10(6) 前胸背板在中央縱沟每旁有盘部前陷及盘部后陷各一.....                  | <i>Asiobates</i>       |

### 亚屬 *Henicocerus* Steph.

本亚屬只發現一种，为 *Ochthebius salebrosus* 新种。这个新种的發現，是本亚屬昆虫在我国的首次紀錄。本亚屬昆虫以前在印度馬來亞区也沒有發現过。

#### *Ochthebius (Henicocerus) salebrosus*, sp. nov.

体长 1.48 毫米，寬 0.68 毫米。体色黑，形长，背面極粗糙不平。

头部色黑，額头顶区連复眼成梯形，中部隆起，隆起部略呈长三角形，其上有十多个頗深刻点，刻点間較光滑，一般比刻点小，前部無刻点，質地粗糙，隆起部每旁有一大而呈瓜仁形的凹陷，凹陷大小約与中央隆起部相等，凹陷区質地粗糙，顏色黑而带棕。口上縫頗深而弯。唇基呈矩形，稍隆起，刻点頗大，刻点間隙光滑，一般較刻点窄。上唇比唇基大，略呈半圆形，有頗深的刻点，刻点間隙光滑，前部無显著刻点，前緣中部有深缺切。下顎鬚色較淡，末前节长大，末节微小，末端平截。顏棕黑色，光耀，近于矩形，有刻点，刻点大小不規則，前緣和側緣几乎平直。

前胸背板黑色，稍呈心形，但寬度仍稍超过长度。盘状部略呈六角形，刻点大而深，密，刻点間隙光滑，一般比刻点窄，但側部的間隙稍寬。中央縱沟中部狹窄。盘状部有盘部前陷及盘部后陷各一对，盘部前陷的界綫頗清楚，但盘部后陷的界綫不大清楚，因而，盘状部后部現出了一条不显著的橫陷。在盘部后陷的外方，还有一个凹陷，其位置近前胸背板后角。上述三对凹陷大小相差不远，只是盘部后陷界綫不大清楚，其大小頗不易观察。盘状部与耳状部之間無显明分沟，整个盘状部隆起，其側緣頗陡直，下方外側即为耳状部。

耳状部色較淡，光滑，其寬度約与盘状部的前緣的相等，有粗糙而深色的邊緣。前胸背板前緣近于平直；側緣由中部稍后处縮窄，縮窄部有窄透明膜圍繞着；后緣稍后突。前胸背板前角圓，后角近直角形。

小盾片色黑，小，三角形。

胸部腹板色黑，前胸腹板滿布微小顆粒，有一橫沟。中胸腹板光耀，后部后伸于中足基节間。后胸腹板粗糙，側部披毛頗密。

鞘翅色黑，长度約等于前胸背板 2.5 倍，刻点排成縱列点紋，刻点頗大，刻点之間的部分显著地隆起，有許多刻点由小橫沟互相連通，結果形成了在鞘翅上有許多不規則的橫沟形象。縱列点紋的間条較刻点窄，有許多很微小的縱形隆起或粒状隆起。由于上述各种构造关系，鞘翅显出了凹突不一、高低不平的形象。鞘翅側緣稍弯拱，突緣显著，前角圓，后角寬圓。

足棕黑色。前足基节稍呈球形，薄披短毛。中足基节柄状，質地与前足同。腿节光耀，有微小而稀疏的刻点。脛节短，有頗稀疏的刺。跗节末节大而长，长度約与前 4 节的长度相等。爪纖細而弯。

腹部基部 5 腹板暗黑色，粗糙，披柔毛。第 6 腹板光耀，薄披稀疏的小毛。第 7 腹板黑色，小，刻点均匀，后部有一不明显的橫沟。

完模标本：♀，著者于 1939 年 11 月 12 日采自云南澂江。采集地点为一終年积水水池，其中水生植物頗丰富。完模标本存中山大学昆虫标本室。在云南澂江这一种是十分稀少的，著者在該地区經 7、8 个月不断采集，只采得标本 1 头。

本种与 *O. exsculptus* Germ. 及 *O. halbherri* Reit. 的区别在于額头顶区的刻較深大和該区的光滑質地。

### 亚屬 *Aulacochthebius* Kuw.

本亚屬的昆虫只發現一种，为 *Ochthebius hunanensis* 新种。这个新种的發現，是本亚屬昆虫在我国的首次紀錄。

#### *Ochthebius (Aulacochthebius) hunanensis*, sp. nov.

体长 1.00 毫米，寬 0.50 毫米。体色棕黑，体形长。

头部黑色，光滑，無显著刻点或刻紋，披頗长而平伏的蓝色毛，毛易脫落，制成的标本，可能脫去許多毛。額头顶区橫形，色稍带棕，中部有一长形而光滑的隆起部，側部有近于三角形的大形凹陷，凹陷的前后部均隆起，前部隆起較大。复眼突出。口上縫深，稍弯拱。唇基色黑，光滑，中部稍隆起，無显著刻点或刻紋。上唇粗糙，比唇基小，近半圓形。下顎鬚短，棕黑色，第 3 节特大而长，端部加厚，第 4 节小，末端尖細。頰黑色，矩形，刻点頗深大，刻点間隙光滑，一般比刻点窄。

前胸背板極寬，光滑，棕黑色，披頗长而平伏的蓝色毛，無显著刻点。盘状部寬度約等于耳状部寬度的三倍。盘状部前部有一深而稍弯的黑色橫沟，該橫沟的每端外側有一长形的凹陷，似橫沟的延伸部；盘状部后部也有一橫沟，长度和顏色与前部橫沟無大区别，但較直，該橫沟的每端外側也有一凹陷；盘状部中央有一縱沟，縱沟較橫沟窄。耳状部質地

与盘状部同,两者由一縱沟分界。耳状部前部也有不大明显的橫沟。盘状部前緣中部些微突出,后緣平直。耳状部前緣成一頗深缺刻,側緣弯缺,后緣近平直,前角突出成指状,后角近直角形。盘状部后角尖突。

小盾片黑色,光耀,三角形。

前胸腹板黑色,粗糙,中央有小縱隆脊,后部后伸于前足基节間,成短刺状。中胸腹板黑色,粗糙,后部后伸于中足基节間成三角形片。后胸腹板黑色,粗糙,披毛,中央部分稍突起,菱形,光耀,其中部有一些小刻点,側部有沟。



圖 1 *O. hunanensis*, 新种, 雄性外生殖器

鞘翅棕黑色,长度約等于前胸背板的 2.5 倍,有極小而疏的毛,有由刻点排成的縱列点紋,这些刻点頗深而大,第 2 至第 4 的縱列点紋中,有两条不达于鞘翅末端,縱紋的間隙光滑,比刻点小,些微隆起。鞘翅突緣不显著,內緣有隆起而光耀的边緣,前角圓,后角寬圓。

足棕黑色。前足基节粗糙,近圓筒形,薄披柔毛。中足基节極粗糙,近柄状。各足腿节光耀,有一些刻点。脛节端部涨大,刺細长,頗密。跗节 5 节,末节最长。爪纖細而弯。

腹部基部 4 腹板暗棕黑色,薄披短毛。第 5 腹板色較浅,光耀,披毛較少。第 6 腹板較长,顏色及質地与第 5 节同,披毛稀疏,第 7 腹板小,光耀,后緣呈半圓形,披毛稀疏,腹部末端有长毛。

雄性外生殖器近于圓筒形。基叶略弯,近中部处有一小缺,缺刻末端的部分漸形尖細。端叶呈膜質,近筒状,端部圓形,有些标本端部卷繞不伸。側叶細长,长度与陽莖約相等,近端部处略胀大,披極短疏毛,端部尖。

完模标本: ♂, 著者于 1941 年 7 月 29 日采自湖南省宜章县的栗源堡村。异模标本: ♀, 采集時間地点同完模标本。副模标本: 29 头, 采集時間地点同完模标本; 5 头, 著者于 1941 年 9 月 5 日采自衡陽; 107 头, 陈湘芷于 1944 年春間采自湖南省攸县。上述标本, 存中山大学昆虫标本室。

本种的个体大小及顏色变异頗大。采自衡陽的标本, 体色較黑, 体积較大。采自宜章和攸县的标本, 体軀較小, 体色也較淡。

本种在湖南省宜章县一带是一个常見种类, 自春日开始在水稻田、灌溉水渠、积水坑等处很容易采到, 在冬天个体数量减少。

本种近似 *O. asiaticus* Orchym., 但与該种有下列区别: 前胸背板的耳状部側緣弯陷不深, 第 1 至第 4 腹节腹板的毛不浓密, 雌虫雄虫第 6 腹节的腹板比第 5 腹板长。*O. asiaticus* Orchym. 的原叙述十分簡單, 又未描述雄性外生殖器, 但根据上列几点形态上的不同, 本种和該种有種的区别。

### 亚屬 *Asiobates* Thoms.

本亚屬的昆虫, 在我国只發現一种, 为 *Ochthebius nitidus* 新种。这个新种的發現, 是本亚屬昆虫在我国的首次記錄。

*Ochthebius (Asiobates) nitidus*, sp. nov.

体长 1.92 毫米, 宽 0.84 毫米。体色棕黑, 体形长。

头部棕黑色。额头顶区有稀疏而不规则的刻点, 刻点间隙光滑; 额头顶区的中部近前缘处有一长形小凹陷, 接口上缝, 后部有一近于菱形的小凹陷; 额头顶区在复眼间有两个大形凹陷, 凹陷的大小因个体不同而有变异。口上缝宽, 两侧尤宽, 稍弯拱。唇基中部隆起, 有稀疏小刻点, 刻点间光滑。上唇有不规则刻点, 刻点间隙光滑, 前缘稍弯陷。下颚鬚色较头部色浅, 末前节大, 棒状, 末端色较深, 末节末端尖。颜棕黑色, 近矩形, 有稀疏及规则的刻点, 刻点间隙光滑, 前缘弯陷, 侧缘平直。

前胸背板棕黑色, 但颜色较头部色浅, 有一些个体前胸背板的颜色和头部颜色没有显著的差别。前胸背板宽度显著地超过长度, 盘状部有不规则的刻点, 刻点比头部背面的刻点大得多, 刻点间隙光滑。中央纵沟颇深, 两端尖细。盘部前陷深, 斜置, 近于椭圆形, 盘部后陷深, 较前陷大, 形长而斜置。盘状部与耳状部之间的沟深。耳状部外侧及后部有刻点。前胸背板前缘有狭窄膜质缘, 中部稍突出, 两侧各有一不显著的小弯陷; 侧缘由后方三分之一处起缩窄, 缩窄处起点有一小尖突, 缩窄部有长形透明膜围绕着, 这透明膜绕过了前胸背板的后角; 后缘稍后突。前胸背板前角宽圆, 后角近于圆形。



圖 2 *O. nitidus*, 新种, 雄性外生殖器

小盾片棕黑色, 三角形, 粗糙, 端部光滑。

胸部腹板棕黑色, 粗糙。前胸腹板后伸于前足基节间, 中央有纵形隆脊。中胸腹板后部后伸于中足基节间, 成三角形小片。后胸腹板两侧披毛, 中部光滑。

鞘翅棕黑色, 长度约等于前胸背板 3 倍, 披有极小的毛, 有由颇大的刻点排列而成的纵纹, 纵纹的间条微隆起, 无刻点, 较刻点稍宽。鞘翅基端近内缘处稍隆起, 鞘翅前部三分之一处有一不明显的大形浅陷, 缘突显著, 前角圆, 后角宽圆。

足颜色较背面稍浅。前足及中足基节柄状, 披毛。各足腿节有不规则刻点, 披薄毛。各足胫节稍呈棒状, 披密而粗的刚毛。跗节末节长, 长度约等于前数节相加长度。爪纤细而弯。

腹部基部 5 腹板暗黑色, 粗糙, 披厚毛。第 6 腹板色较淡, 光耀, 其后部有细小刻点及稀疏短毛。第 7 腹板颜色与第 6 节相同, 后部有长毛。

雄性外生殖器的基叶长而弯, 基端前方缩窄, 基端呈半球状。端叶短, 呈钟状, 末端有陷, 陷内有一小棒。侧叶细长, 端部有一些短毛。

完模标本: ♂, 著者于 1940 年 3 月 26 日采自云南省禄丰村的一条溪水中。异模标本: ♀, 采集日期地点与完模标本同。副模标本: 4 头, 采集日期及地点与完模标本同。上述标本存中山大学昆虫标本室。

本种与 *O. amplipollis* Champ. 及 *O. kosiensis* Champ. 的区别在于背面没有明显的毛; 和 *O. laevisculptus* Reit. 的区别在于鞘翅有较大的刻点; 和 *O. atricapillus* Reit. 的

区别在于上唇前缘的弯陷浅而平滑。

### 亚属 *Homalochthebius* Kuw.

本亚属昆虫,在我国过去曾有一种的纪录即 *O. yunnanensis* Orchym., 其后,著者在我国发现新种一种及新亚种一种。现将各种制成检索表,并分别加以叙述。

#### *Homalochthebius* 亚属的种及亚种检索表

- 1(2) 鞘翅有不规则黑色斑..... *unimaculatus*, sp. nov.
- 2(1) 鞘翅无明显黑斑
- 3(4) 上唇有许多微小颗粒状突起,质地粗糙。唇基比额头顶区小..... *yunnanensis* Orchym.
- 4(3) 上唇无颗粒状突起。唇基与额头顶区几乎等大..... *yunnanensis furcatus*, subsp. nov.

#### *Ochthebius* (*Homalochthebius*) *unimaculatus*, sp. nov.

体长 2.20 毫米,体宽 0.90 毫米。体色棕黑,体形长卵形。

头棕黑色,有一些个体呈黑色,也有一些个体的头部背面微带紫铜色(重庆标本),有些个体带青铜色(曲江标本),头部一般无明显短毛,但曲江标本有较明显的短毛。额头顶区的刻点在完模标本和异模标本是相当密的,重庆的一些标本刻点较疏,曲江标本的刻点很不规则,在中部有一些刻点连成短刻纹,使该部形成粗糙凹突的质地,各地标本的刻点间隙均光滑。复眼之间有两个大形凹陷,每凹陷中部有裂条状纵纹。上唇和额头顶区约等大,有刻点,雄虫上唇的前缘稍弯陷,雌虫的弯陷较深。下颚鬚色较头部浅,末前节大,棒状,末节小,末端尖细。上颚外侧有短而颇粗的毛列。额棕黑色,稍呈矩形,有颇疏刻点,刻点间隙光滑,额的前缘稍弯陷。



图 3 *O. unimaculatus*, 新种,  
雄性外生殖器

前胸背板颜色与头部的相似,宽度显著地超过长度。盘状部刻点颇深大,中部的刻点较疏,后部的较密,前部的较小,有一些个体后部的刻点也较小。中央纵沟窄,前部尖,前后两端均不达背板的前后缘,雌虫的中央纵沟较宽,但采自曲江的雄虫标本的中央纵沟也较宽。耳状部和盘状部之间的凹陷大,耳状部有刻点,刻点和盘状部的差不多相同。前胸背板前缘稍弯陷,绕以透明的窄膜;侧缘由后部三分之一处缩窄,缩窄部有透明的膜围绕,这透明膜绕过了前胸背板的后角,在侧缘开始缩窄的地方,有一个微小的刺状突起;后缘中部稍后突。前胸背板前角圆,后角近直角形。

小盾片黑色,三角形,基部色暗而粗糙,端部光滑。

胸部腹板色黑,粗糙。中胸腹板披毛,后部呈三角形,三角形部分的后端呈刺状,后伸于中足基节间。后胸腹板披厚毛,后部中央有一光滑无毛的小区。

鞘翅色较前胸背板的稍浅,长度约等于前胸背板 2.5 倍,每鞘翅有一个稍呈弯形而不大明显的黑斑,该黑斑位于鞘翅基部,占鞘翅一半以上的长度,在鞘翅颜色较浅的标本,这个黑斑较为显著。每鞘翅有 10 条深显的纵列点纹,此外,有由一点或两点所成的小盾片纹,有一些标本的小盾片纹不显著。鞘翅纵列点纹的间条光滑,稍隆起,约与点纹等宽。

鞘翅緣突頗显著，側緣弯拱，有隆起小邊緣。鞘翅前角圓，后角寬圓。

足棕色。前足基节粗糙，錐狀，薄披短毛。中足基节粗糙，近球形，質地与前足基节同。各足腿节光耀，有微小而不規則的刻点。脛节光耀，有刺。跗节末节长，前足中足末节长度約等于前数节相加长度，后足末节长度稍短。爪纖細而弯。

腹部基部<sup>5</sup>腹板暗黑色，披毛。第6腹板棕色，光耀，基部有微小刻紋，端部有小毛。第7腹板稍呈半圓形，后緣有一列长毛。

雄性外生殖器的基叶长形，微弯，基部前方縮窄。端叶短于陽茎基極多，端部扩伸呈喇叭口狀。側叶細长，长度短于基叶，端部有一些短毛。

完模标本：♂，著者于1940年10月8日采自貴陽。异模标本：♀，采集日期与地点与完模标本同。副模标本：23头，采集日期及地点与完模标本同；1头，林世平于1941年10月22日采自广东省曲江；49头，陈湘芷于1942年3月26日采自重庆。上述标本存中山大学昆虫标本室。

本种雄虫上顎外緣的刚毛比雌虫的粗得多，这特征是本种的主要第二性征。雌虫上唇前緣弯陷較雄虫的深，这也是雌雄分別的一个好特征。

本种和 *O. perudus* Reit. 的區別在于具有小盾片縫；和 *O. coomani* Orchym. 的區別在于前胸背板的刻点較深而大。

### *Ochthebius (Homalochthebius) yunnanensis* Orchym.

*Ochthebius (H.) yunnanensis* Orchymont 1925, Ann. & Bull. Soc. Ent. Belg., 65:263; Orchymont 1935, Peking Nat. Hist. Bull., 9:188; Wu 1936, Cat. Ins. Sin., 3:271; Orchymont 1942, Bull. Mus. Hist. Nat. Belg. 18(39):16.

1925年 d'Orchymont 叙述了这一个种。标本采自云南，可是沒有小地名，也有几个标本的标签写着采自“云南或日本”。模式标本存 d'Orchymont 处和柏林昆虫博物館。1942年 d'Orchymont 补充描叙并圖示这一种昆虫的雄性外生殖器。著者于1939及1940年在云南各地采得这个亚屬的标本凡数百头，采集地点包括昆明、昆陽、晋宁、大理、潞江等县，标本外形均極相似。昆明、昆陽、晋宁和大理的标本，其雄性外生殖器的构造和 *O. yunnanensis* Orchym. 的相同，潞江标本的雄性外生殖器就显著地不同，其他构造也有不同处，因此，潞江的种类是有別于 *O. yunnanensis* 的，現鑒定为一新亚种，叙述見后。

### *Ochthebius (Homalochthebius) yunnanensis furcatus*, subsp. nov.

体长2.10毫米，寬0.90毫米。体形长，鞘翅近长卵形，体色棕褐。

头部棕黑色，有金屬光澤，披稀疏短毛。額头顶区有深而較大的刻点，刻点間隙光耀，較刻点微寬，有些标本的刻点連成許多不規則的刻紋；复眼間有两个大形凹陷，凹陷中央有縱形裂沟，凹陷外側有一个光滑的部分。口上縫寬，拱形。唇基薄披短毛，側緣直，刻点較疏而較小，無显著的顆粒狀小突起(雌虫的有一些)。上唇大，差不多与唇基等大，刻点的情况相似，但稍疏，沒有顆粒狀小突起，前緣仅稍弯缺，側緣圓形。下顎鬚棕紅色，末前节涨大，末节末端尖細。顏棕色，近距形，前緣近于平直，有疏而頗大的刻点，刻点間隙光滑。

前胸背板棕色, 宽度显著地超过长度。色较头部浅。盘状部中央色黑, 披不显著的微小短毛, 有深刻点, 刻点比额头顶区的大而密, 刻点间隙有些比刻点宽, 有些比刻点窄。盘状部中沟窄, 黑色, 前后端均不达鞘翅前后缘, 中部较窄, 前部较宽, 有些标本中沟的两端宽度和中部差不多一样。中沟后部有一色较前胸背板稍深而不甚明显的横形浅陷。盘状部和耳状部分界的沟宽大而稍弯, 耳状部色较盘状部浅, 刻点和盘状部相似, 但较浅较小也较疏。前胸背板前缘颇平直, 有窄膜质缘; 侧缘前部四分之三圆拱, 后部自三分之一处起缩窄, 缩窄处有透明膜围绕, 这透明膜也绕过后角; 后缘稍后突。前胸背板前角圆, 后角直角形。



图 4 *O. yunnanensis furcatus*, 新亚种,  
雄性外生殖器

小盾片色黑, 三角形, 基部有微小刻纹, 其他部分光滑。

胸部腹板色黑, 粗糙, 披薄毛。前胸腹板横形, 后部中央后伸于前足基节间, 呈三角形。中胸腹板中央有由小隆脊包围而成的杏仁形部分, 后部后伸于中足基节间。后胸腹板大, 后部中央有一近心形的光滑部分。

鞘翅颜色与前胸背板的相似, 长度约等于前胸背板 3 倍, 披稀疏短毛。鞘翅刻点纵行列深而大, 小盾片缝短, 有些标本的小盾片缝只由一点所成, 纵纹的间条光滑, 稍隆起, 宽度约与纵纹等大或稍大。鞘翅缘突颇显著, 前角圆, 后角宽圆。

足棕色。前足基节近圆锥形, 粗糙, 端部披毛。中足基节长球形, 粗糙, 端部披毛较前足密。各足腿节有疏而带毛的微小刻点。各足胫节有粗密的刚毛。前足及后足的跗节末节长度与基部数节相加长度约相等, 中足的略短于基部数节相加长度。不纤细。

腹部基部 5 腹板棕黑色, 粗糙, 披毛。第 6 腹板棕色, 光耀, 雄虫披稀毛, 后缘稍弯陷, 第 6 腹板基部有微小刻纹。雄虫第 7 腹板小, 近半圆形, 后缘有毛, 雌虫第 7 腹板小, 后部中央稍突, 两旁平直。

雄性外生殖器的基叶长而弯, 基端前方缩窄。端叶短, 端部有一个钳状的构造, 两个钳支的大小不同。侧叶细长, 较基叶短, 端部有一些短毛。

完模标本: ♂, 著者于 1939 年 4 月 26 日采自云南省潞江县。异模标本: ♀, 采集日期地点与完模标本同。副模标本: 140 头, 采集日期地点与完模标本同; 60 头, 于 1939 年 9 月采自同上地点; 20 头, 于 1940 年 1 月采自同上地点。上述标本存中山大学昆虫标本室。

本亚种极似 *O. yunnanensis* Orchym., 但体色一般较深, 唇基较大, 且无明显的颗粒状小突起, 上唇也没有这种突起。*O. yunnanensis* Orchym. 的唇基和上唇都有较明显的颗粒状突起。

### 亚属 *Bothochius* Rey

本亚属的昆虫, 在我国前后发现 5 个种。现知的种类, 分布于新疆、陕西、云南、西藏等地。



***Bothochius* 亚属的种检索表**

- 1(6) 上唇前缘弯陷深
- 2(3) 背面有微小颗粒, 质地粗糙..... *verrucosus* Pu
- 3(2) 背面质地光滑
- 4(5) 鞘翅的刻点有一些排列成行..... *lobatus*, sp. nov.
- 5(4) 鞘翅的刻点不排列成行..... *turkestanus* Kuw.
- 6(1) 上唇稍弯陷或几乎平直
- 7(8) 前胸背板稍呈心形..... *caucasicus* Kuw.
- 8(7) 前胸背板较横置..... *flexus*, sp. nov.

***Ochthebius (Bothochius) flexus*, sp. nov.**

体长 1.82 毫米, 宽 0.84 毫米。体色棕黑, 背面有平伏蓝色毛, 体形长。

头部棕黑色, 有铜色光泽。额头顶区有小而颇深的刻点, 但有些个体的刻点不大明显, 刻点间隙光耀, 一般比刻点宽。额头顶区在复眼之间有一对大而深的凹陷, 其后有一小形的凹陷。口上缝宽, 弯拱。唇基的刻点和额头顶区的相似。上唇大, 粗糙, 有刻点及毛, 呈半圆形, 前缘几乎平直。下颚鬚短, 色较头部背面的浅, 末前节大而长, 末节末端尖细。颧色较头部背面的浅, 近梯形, 有疏而不规则的刻点, 前缘弯陷, 侧缘稍弯陷, 后缘平直。

前胸背板颜色与头部相同, 宽度显著地超过长度。盘状部刻点的大小及深度与额头顶区的相似, 近前缘及近后缘的刻点较密, 刻点间隙光耀, 比刻点宽。背板中央纵沟颇深, 前部扩大, 末端尖细, 有些标本前部不扩大, 有些标本的中沟两端都扩大。盘部前陷及盘部后陷近椭圆形, 前陷较短, 后陷较长, 稍斜向, 有些标本的前陷较圆, 后陷较短较浅。盘状部和耳状部之间的沟长形, 大而颇深。前胸背板各凹陷的质地均粗糙。耳状部的占前胸背板长度三分之一, 颜色较浅, 质地凹陷不平, 耳状部侧缘几平直。前胸背板前缘微弯陷, 有狭窄的膜质缘; 侧缘由前部三分之一处向内缩窄, 缩窄部稍弯, 并绕以膜质缘, 该膜质缘绕过前胸背板后角; 后缘中部突出。前胸背板前角及后角圆宽。

小盾片色黑, 三角形, 光滑。

胸部腹板暗黑色, 粗糙, 披毛。前胸腹板中央稍隆起成纵形隆脊, 后部后伸于前足基节间成刺状。中胸腹板的小隆脊围成三角形, 后部后伸于中足基节间成长三角形的刺状构造。后胸腹板大, 披毛较前胸、中胸的厚。

鞘翅棕黑色, 色较胸稍淡, 长度约等于前胸背板 3 倍, 刻点颇深而大, 有一些刻点排成纵列, 刻点间条光耀, 一般比刻点大, 有极微小而密致的纹, 雌虫比较明显。鞘翅的内外缘均有小边缘, 内缘的小边缘以后部的较明显。鞘翅的缘突及侧沟颇明显, 前角颇锐, 后角圆宽。

足棕黑色, 较体躯腹面颜色浅。前足基节柄状, 粗糙, 披毛。中足基节近球形, 粗糙, 披毛。各足腿节有披毛的疏刻点。各足胫节近直形, 端部稍大, 前足胫节刚毛短, 中足胫节刚毛较长而密, 后足胫节刚毛长而疏, 各胫节近端部的毛均较长及较粗。跗节末节长,




圖 5 *O. flexus*, 新种, 雄性外生殖器

前三节的末端各有一小毛。爪纖細而弯。

腹板基部<sup>5</sup>腹板暗黑色,橫形而窄,粗糙,披毛。第6腹板棕黑色,窄,基部粗糙,端半部有微小刻点及纖小的毛。第7腹板棕黑色,光耀,有微小刻点,后緣弯拱,中部有微小缺刻。

雄性外生殖器的基叶长,極弯曲,基部前縮窄,中部極狹小,末端尖。端叶比基叶短極多,末端圓拱。側叶細长,末端有短毛。

完模标本:♂,著者于1940年3月26日采自云南省陆丰村。异模标本:♀,采集日期地点与完模标本同。副模标本:7头,采集地点与完模标本同。上列标本存中山大学昆虫标本室。

本种与 *O. turkestanus* Kuw, *O. nobilis* Villa 及 *O. rugusae* Kuw. 的区别在于几乎平直的上唇前緣;和 *O. detritus* Rey 的区别在于前胸背板前緣弯陷極微;和 *O. caucasicus* Kuw. 的区别在于較為橫形的前胸背板。

### *Ochthebius (Bothochius) lobatus*, sp. nov.

体长2.16毫米,寬1.08毫米。

本种極似 *O. flexus*, 与該种之区别在于下列特征:体积較大,头部及前胸背板的刻点較不清楚,上唇前緣的缺刻較深,鞘翅無密致的小紋而有一些較疏而微小的长形裂紋狀刻紋。本种雄性外生殖器基叶在近基部三分之一处縮窄,基叶末端較尖长,端叶較圓。*O. flexus* 的雄性外生殖器基叶在近中部处縮窄,末端胀大部分比 *O. lobatus* 的大,形状也不同;端叶近长筒形。基于上述主要特征,本种可与 *O. flexus* 有種的区别特征。



圖6 *O. lobatus*, 新种, 雄性外生殖器

完模标本:♂,陈湘芷于1942年8月8日采自四川重庆。异模标本:♀,采集時間与地址与完模标本同。副模标本:26头,与完模标本同时同地采集。上述标本存中山大学昆虫标本室。

### 亚屬 *Hymenodes* Muls.

本亚屬昆虫,我国过去未有紀載,本文紀录2种,均新种。

#### 亚屬 *Hymenodes* 的种检索表

- 1(2) 前胸背板中央縱沟完整.....*exilis*, sp. nov.
- 2(1) 前胸背板中央縱沟中断,分为前后二部.....*octofoveatus*, sp. nov.

### *Ochthebius (Hymenodes) exilis*, sp. nov.

体长1.80毫米,寬0.80毫米。体色深棕黑,有金屬光澤,背面有平伏的长毛。

头深棕黑色,刻点大,疏,不大清楚。額头頂区及唇基有微小的顆粒突起,致質地粗糙;額头頂区在复眼間有两个近橢圓形的深陷,其后有一較小凹陷。口上縫稍弯拱。唇基

近梯形。上唇半圆形,略小于唇基,有密而颇粗的刻点,披毛颇长而密。下颚鬚棕色,末前节大而长,末节端部稍尖细。颊棕黑色,近四角形,有微小颗粒,并有规则刻纹及刻点,前缘中部弯陷颇深,侧缘稍弯陷。

前胸背板色如头部,宽度比长度稍大,盘状部有极微小颗粒突起及颇疏而小的刻点,中央纵沟后部缩窄,前后两端尖细,其后端几乎达到背板后缘,前端不达前缘。盘部前陷近圆形,盘部后陷长椭圆形。盘状部和耳状部分界的沟大,稍斜置,其前端不达前缘。耳状部也有颇疏的小刻点。前胸背板前缘近平直;侧缘在中部之前以后部分向内缩窄,缩窄部有透明薄膜,该薄膜围绕前胸背板后角;后缘向后突。前胸背板前角近直角形,后角圆。



图 7 *O. exilis*, 新种, 雄性外生殖器

小盾片深棕黑色,三角形,光耀。

胸部腹板色黑,粗糙。前胸背板披柔毛,中部稍成脊状隆起,后部后伸于前足基节间成三角形突起。中胸腹板侧缘披毛,中部隆起成长三角形片,并后伸于中足基节间。后胸腹板厚披白毛,后缘中部有一小凹陷。

鞘翅深棕黑色,色较头部及胸部的稍浅,长度约等于前胸背板 2.5 倍,每鞘翅有 10 行刻点,刻点间条稍隆起,有微小颗粒突起,宽度与刻点略等稍大。鞘翅内缘有小光滑边缘,外缘稍弯拱,有隆起小边缘。鞘翅缘突显著,前角直角形,后角宽圆。

足棕黑色,前足及中足基节近柄状,粗糙,披毛。腿节有不规则的带毛刻点。胫节端部大,有粗大而颇密的刺毛。跗节末节长,约与前数节等长。爪纤细而弯。

腹部基部 5 腹板暗黑,窄,粗糙,披毛。第 6 腹板较宽,色较浅,光耀,粗糙,有些标本该节基部不粗糙。第 7 腹板小,光耀,近半圆形。腹部末端有一些纤细刚毛。

雄性外生殖器的基叶长而稍弯,略向端部尖细基端前方缩窄。端叶短,近刀片状。侧叶细长,其端部有短毛。

完模标本: ♂, 1940 年 3 月 20 日著者采自云南省陆丰村。异模标本: ♀, 采集时间地点与完模标本同。副模标本: 8 枚, 采集时间地点与完模标本同。上述标本存中山大学昆虫标本室。

本种雌雄之别在于雌虫腹部末端无显著刚毛。

本种与 *O. foveolatus* Germ., *O. nitidipennis* Champ., *O. flumineus* Orchym. 及 *O. explanatus* Orchym. 的区别在于上唇前缘无显著弯陷; 与 *O. cupricollis* Champ. 之别在于较长的鞘翅; 与 *O. scintillans* Champ. 及 *O. sexfoveatus* Champ. 之别在于鞘翅刻间条有微小颗粒突起; 与 *O. strigosus* Champ. 之别在于前胸背板只有 5 个凹陷。

### *Ochthebius (Hymenodes) octofoveatus*, sp. nov.

体长 1.68 毫米, 宽 0.80 毫米。体深棕色, 有铜色光泽。

头部深棕色, 有微小颗粒状突起。额头顶区带红铜色光泽, 有少数刻点, 复眼间有两个大形凹陷, 其后有一较小凹陷, 每个大形凹陷的后侧方有一小形突起部, 额头顶区中央有一近长方形突起部分, 该突起部后方有一浅纵陷。唇基横形, 色较额头顶区深。上唇色

略浅于唇基，长度约与唇基相等，有不显著的微小颗粒状突起，有刻点，前缘中部有呈半圆形的深缺刻。下颚鬚深棕色，末前节大，近锤状，末节末端尖细。颜棕色，近四边形，粗糙，有微小而不规则的刻点，前缘稍弯陷。

前胸背板色似唇基，宽度较长度略大。盘状部有似头部的微小颗粒状突起。中央纵沟两端尖细，均不达前胸背板的前后缘，该纵沟分为前后二部，前部的沟略长于后部。盘部前陷近椭圆形，盘部后陷长形，稍斜向，前后陷的宽度与中沟约相等。盘部前陷与后陷的两旁均各有一凹陷，在前陷两端者较小，在后陷两端者较大。盘状部及耳状部的分界沟长而大。耳状部色浅于盘状部，也有微小颗粒状突起。前胸背板前缘几乎平直，有窄膜缘；侧缘由前部三分之一处向内方缩窄，缩窄部有薄膜围绕，该薄膜绕过前胸背板后角；后缘有窄膜缘，中部后突；前胸背板前角近直角形，后角圆宽。

小盾片三角形，有极微小颗粒状突起。

前胸及中胸腹板棕黑色，有颗粒状突起。前胸腹板中央有隆脊，后部后伸于前足基节间。中胸腹板后部成三角形片，后伸于中足基节间。后胸腹板大，棕色，粗糙，薄披柔毛，后缘中部有小缺刻。

鞘翅色较前胸背板浅，长度约等于前胸背板 2 倍，每鞘翅有 10 行刻点，刻点间条光滑，略隆起，一般比刻点宽或与刻点等宽。鞘翅内缘有一边线，侧缘弯拱，有隆起小缘。鞘翅缘突显著，前角圆，后角圆宽。

足棕色。前足及中足的基节近柄状，粗糙，薄披柔毛。各足腿节和胫节有微小刻纹。胫节的刺毛颇密。跗节末节长而大。爪纤细而长。

腹部腹板棕色，粗糙，基部 5 腹板窄，薄披柔毛。第 6 腹板较宽，中部光耀，有微小刻点，基部侧方粗糙有柔毛，端部侧方有不规则小刻点及柔毛。第 7 腹板近半圆形，光耀，有不规则的微小刻点，基部有不显著的刻纹。

完模标本：♀，著者于 1940 年 3 月 26 日采自云南省陆丰村，存中山大学昆虫标本室。

本种与 *O. foveatus* Germ. 及 *O. nitidipennis* Champ. 之别在于鞘翅的刻点行列的间条稍隆起；与 *O. cupricollis* Champ. 的区别在于鞘翅较长；与 *O. sexfoveatus* Champ. 之别在于鞘翅没有显著的柔毛；与 *O. exilis* 新种、*O. flumineus* Orchym. 及 *O. explanatus* Orchym. 之别在于前胸背板中央纵沟分成前后两部分。

## THE GENUS *OCHTHEBIUS* LEACH OF CHINA (COLEOPTERA, PALPICORNIA)

PU CHIH-LUNG

Department of Biology, Sun Yatsen University

The present paper deals with twelve species and one subspecies of the genus *Ochthebius* from China, of which eight species and one subspecies are new to science.

### *Ochthebius (Henicocerus) salebrosus*, sp. nov.

Size, 1.48 mm. long, 0.68 mm. wide.

Head significantly narrower than pronotum; fronto-vertex black, with moderately

large and dense punctures, intervals glabrous, narrower than the punctures and generally raised into ridges, two large irregular depressions between the compound eyes; epistomal suture arched; clypeus coarsely and irregularly punctured, intervals glabrous and slightly raised; labrum larger than the clypeus, proximal area with transversely arranged punctures; intervals glabrous, slightly raised, anterior margin semicircular, deeply emarginate medially; maxillary palpus lighter in color, penultimate segment large and long, ultimate segment minute, distal extremity truncate; mentum dark brown, transverse, shining, somewhat quadrangular, nearly straight anteriorly and laterally; punctures irregular in size.

Pronotum black, more or less heart-shaped; disc coarsely and densely punctured, intervals glabrous and generally narrower than the punctures; median longitudinal furrow deep, constricted medially, anterior and posterior depressions moderately deep, a depression situated exteriorly to the posterior depression; outer depressions large; anterior margin nearly straight, lateral margin retracted posteriorly from a point just behind the middle, the hind part of retracted portion lined with a narrow translucent membrane, posterior margin slightly arched; anterior angles rounded, posterior angles rectangular.

Scutellum black, triangular.

Prosternum dark brown, large, granulate, with a transverse groove; mesosternum black, shining, hind area extended posteriorly between the middle coxae; metasternum dark brown, roughened, much larger, thickly pubescent laterally.

Elytra black, punctures of longitudinal series moderately large; intervals narrower than the punctures with minute protuberance, the punctures of each series communicated with those of the others by transverse minute grooves; interstices glabrous, very narrow and raised into ridges; lateral margin slightly arched; humerales moderately prominent; anterior angles rounded, posterior angles broadly rounded.

Legs brown; front coxae more or less globular, thinly pubescent, middle coxae somewhat peg-like, with the same texture; femora shining, minutely and sparsely punctured; tibiae with short and moderately sparse spines; distal segment of tarsi large and long, subequal to the preceding segments together; claws slender and curved.

Five proximal abdominal sterna dull black, roughened, pubescent; sixth sternum shining, sparsely and minutely pubescent, median area minutely punctured; seventh sternum black, smaller, evenly punctured, posterior area with an indistinct transverse groove.

Holotype, one female from Chengkiang, Yunnan. November 12, 1939, collected by the author, in the insect collection of Sun Yatsen University.

***Ochthebius (Aulacochthebius) hunanensis, sp. nov.***

Size, 1.00 mm. long, 0.50 mm. wide.

Head black, glabrous, with long procumbent blue hairs; fronto-vertex transverse, a globular glittering and prominent swelling at the middle, two irregular depressions laterally near the compound eyes; a glittering swelling in front and a smaller one behind each depression; epistomal suture deep, slightly arched; clypeus glabrous; labrum roughened, semicircular; maxillary palpi dark brown, short, third segment large and long, thickened distally, fourth segment tapering distally; mentum black, somewhat rectangular, moderately coarsely and deeply punctured, intervals shining, generally narrower than the punctures.

Pronotum very transverse, dark brown, shining, with long procumbent blue hairs, without apparent punctures; anterior discal transverse groove moderately deep, an elongated depression at each of its extremities, posterior discal transverse groove as deep as the anterior one with an elongated depression at each extremity; median longitudinal furrow moderately deep, narrower than the transverse groove; anterior margin arched; lateral margin lined with narrow translucent membrane, slightly concave anteriorly,

retracted posteriorly; posterior margin arched; anterior angles pointed, posterior angles rectangular.

Scutellum black, shining, triangular.

Prosternum black, roughened and transverse, posterior area prolonged posteriorly to form a carina between the anterior coxae; mesosternum dark brown, roughened, posterior area prolonged posteriorly between the middle coxae; metasternum black, roughened and pubescent, middle area raised into a large, glabrous rhomboid area and with a depressed area laterally and posteriorly to relieve the posterior femur.

Elytra dark brown, with sparse minute hairs and a moderately deep large series of setigerous punctures, intervals and interstices shining, generally narrower than the punctures, very slightly raised; humerales not prominent; inner margin with a glabrous border; anterior angles rounded, posterior angles broadly rounded.

Legs dark brown; anterior coxae roughened, somewhat cylindrical, very thinly pubescent; middle coxae somewhat peg-like, very roughened; femora glabrous, more or less fusiform, with few punctures; tibiae thickened distally and thinly pubescent, subequal to their respective femora, tibial spines slender and moderately dense; distal segment of tarsi long; claws slender and curved.

Abdomen dark brown; four proximal sterna dull, roughened, and thinly pubescent; fifth, sixth and seventh sterna lighter in color, shining, with sparser pubescence.

Basal lobe of phallus large, somewhat cylindrical, dorsal and ventral portions prolonged into minute and long distal prolongations respectively; distal lobe membranous; lateral lobes slender and long, subterminally setose and swollen.

Holotype, one male from Ichang, Hunan Province, July 29, 1941, collected by the author. Allotype, one female with the same locality and date as the holotype. Paratypes, twenty-nine specimens, with the same locality and date as the holotype; five specimens from Hengyang, Hunan Province, September 5, 1941, collected by the author; one hundred and seven from the same province, spring, 1944, collected by Miss S. C. Chen. All types in the insect collection of Sun Yatsen University.

### *Ochthebius (Asiobates) nitidus*, sp. nov.

Size, 1.92 mm long, 0.84 mm. wide.

Form elongated; fronto-vertex dark brown, sparsely and irregularly punctured, intervals glabrous; fronto-vertex with a small anterior and posterior depression, two large depressions between the compound eyes and a small triangular median depression behind; epistomal suture slightly arched; clypeus elevated medially with sparse minute punctures; labrum dark brown, coarsely punctured, intervals glabrous, shallowly and smoothly emarginate anteriorly; maxillary palpi brown, penultimate segment large, club-shaped, with distal end darker in color, ultimate segment tapering distally; mentum dark brown, more or less rectangular, sparsely punctured, intervals glabrous, anterior margin emarginate, lateral margin straight.

Pronotum dark brown; disc irregularly and moderately coarsely punctured, intervals glabrous, median longitudinal furrow moderately deep with tapering extremities; anterior discal depressions deep and oblique, somewhat elliptical in outline, posterior discal depressions deep, elongated and oblique, outer depressions large; anterior margin slightly arched medially with narrow membranous border; lateral margin retracted from the posterior third and lined with elongated translucent membrane, the latter encircling the posterior angle of pronotum; posterior margin arched; anterior angles broadly rounded, posterior angles less so.

Scutellum dark brown, roughened, triangular.

Thoracic sterna dark brown, roughened; prosternum longitudinally carinate medially; mesosternum extended posteriorly between the middle coxae; metasternum thinly pube-

scent laterally, glabrous medially.

Elytra dark brown, minutely pubescent, with series of moderately large punctures, interstices slightly raised, glabrous, wider than the punctures; lateral margin arched with minute elevated border; inner margin with elevated border; humerales prominent; anterior angles rounded, posterior angles broadly rounded.

Legs dark brown; anterior and middle coxae peg-like, sparsely pubescent; femora somewhat peg-like, irregularly punctured, sparsely pubescent; tibiae somewhat club-shaped, densely spinose; distal segment of tarsi nearly as long as the preceding segments together.

Five proximal abdominal sterna dull black, roughened, densely pubescent; sixth sternum lighter in color, shining, its posterior area finely punctured and sparsely covered with minute hairs; seventh sternum like the preceding in color, but glabrous and with numerous long hairs posteriorly.

Basal lobe of phallus long and curved; distal lobe irregularly triangular, with a hollow at the distal extremity, in which there is a slender rod; lateral lobes slender, much shorter than the basal lobe, with a few minute setae distally.

Holotype, one male from Lou-fong-tsouen, Yunnan Province, March 26, 1940, collected by the author. Allotype, one female with the same locality and date as the holotype. Paratypes, four specimens with the same locality and date as the holotype. All types in the insect collection of Sun Yatsen University.

***Ochthebius (Homalochthebius) unimaculatus*, sp. nov.**

Size, 2.20 mm. long, 0.90 mm. Wide.

Head dark brown or black, some specimens with metallic shining, without apparent pubescence in most specimens; fronto-vertex and clypeus moderately densely punctured in most specimens, intervals shining; labrum large, transverse, punctured, anterior margin slightly emarginated in male, deeply emarginated in female; maxillary palpi brown, large, penultimate segment large, club-shaped, ultimate segment small, tapering distally; the exterior margin of mandible with rows of brownish, short and thick spines, which are thicker in male; mentum brown, middle area black, somewhat rectangular, anterior margin broadly emarginate, moderately sparsely punctured, intervals shining and wider than the punctures.

Pronotum dark brown; disc moderately coarsely punctured, the punctures near the anterior and posterior margins generally somewhat smaller; median longitudinal furrow generally narrower in male with tapering anterior end; outer depressions wide; anterior margin slightly emarginated, with translucent narrow membranous border; lateral margin retracted from the posterior third and lined with narrow translucent membrane which encircles the posterior angle, a minute spine-like outgrowth present at the posterior end of the non-retracted portion; posterior margin slightly arched medially; anterior angles rounded, posterior angles rectangular.

Scutellum black, triangular; proximal area dull and roughened, distal area glabrous.

Thoracic sterna black; prosternum roughened and transverse; mesosternum roughened, pubescent, spine-like posteriorly between the middle coxae; metasternum roughened and heavily pubescent, with a glabrous portion at the posterior area.

Elytra dark brown, with an irregular black mark on each elytron, ten series of circular deep punctures and a very short scutellar suture on each elytron, the punctures separated by narrow intervals, interstices glabrous, slightly raised and subequal to the punctures in width; inner margin with minute black border; lateral margin somewhat arched, with black minute elevated border; humerales moderately prominent; anterior angles rounded, posterior angles broadly rounded.

Legs brown; front coxae roughened, conical, granulated, minutely sculptured and

minutely lightly pubescent; middle coxae roughened, globular, with the same texture as the front coxae; femora brown, shining, minutely and irregularly punctured; tibiae brown, shining, spinose; tarsi with distal segment long, subequal to the preceding segments together on the front and middle legs, but somewhat shorter on the hind legs; claws slender and curved.

Five proximal abdominal sterna dull black and pubescent; sixth sternum brown and shining, its proximal area with minute sculpture, distal area glabrous, minutely pubescent; seventh sternum dark brown, more or less semicircular, posterior margin with a row of long setae.

Basal lobe of phallus elongated, curved; distal lobe somewhat oval in outline, expanded distally; lateral lobes slender, minutely setose distally, shorter than the basal lobe.

Holotype, one male from Kweiyang, Kweichow Province, October 8, 1940, collected by the author. Allotype, one female with the same locality and date as the holotype. Paratypes, twenty-three specimens with the same locality and date as the holotype; forty-nine specimens from Chungking, Szechuen Province, March 26, 1942, collected by Miss S. C. Chen; one specimen from Chiuchiang, Kwangtung Province, October 22, 1941, collected by Mr. Sping Lin. All types in the insect collection of Sun Yatsen University.

***Ochthebius (Homalochthebius) yunnanensis furcatus*, subsp. nov.**

Size, 2.10 mm. long, 0.90 mm. wide.

Head dark brown with metallic shining, thinly pubescent; frontovertex coarsely and deeply punctured, intervals shining, generally broader than the punctures, two large depressions between the compound eyes; epistomal suture arched and broad; clypeus with sparse and small punctures, not significantly granulated; labrum large, with punctures as those on clypeus, but sparser, anterior margin slightly emarginated in the male and deeply notched in the female; maxillary palpi brown red, penultimate segment greatly swollen, ultimate segment tapering distally; mentum brown, sparsely punctured, intervals glabrous, anterior margin nearly straight.

Pronotum brown, disc black, minutely pubescent, coarsely and densely punctured; median longitudinal furrow black, narrow, not reaching the margins of pronotum, an ill-defined transverse depression posteriorly to the median furrow; outer depressions distinct; anterior margin nearly straight; anterior third of the lateral margin arched, posterior fourth retracted, lined with moderately broad translucent membrane which encircles the posterior angle of pronotum; posterior margin slightly arched; anterior angles rounded, posterior angles rectangular.

Scutellum dark brown, triangular.

Thoracic sterna black, roughened and thinly pubescent; prosternum transverse, with posterior area between front coxae triangular; mesosternum with posterior area between the middle coxae spine-like; metasternum with a glabrous heart-shaped area medioposteriorly.

Elytra dark brown, very minutely pubescent; series of punctures deep and large with very short scutellar suture; interstices glabrous, subequal to the punctures in width, slightly raised; intervals of punctures narrow; humerales moderately prominent; anterior angles rounded, posterior angles broadly rounded.

Legs light brown; anterior and middle coxae roughened; femora with sparse setigerous minute punctures and short hairs; tibiae strongly spinose; tarsi with distal segment subequal to the preceding segment together.

Five proximal abdominal sterna dark brown, roughened, pubescent; sixth sternum brown, shining, sparsely pubescent, posterior margin very slightly emarginate; seventh



sternum small, more or less semicircular, pubescent along posterior margin.

Basal lobe of phallus curved, median portion slightly narrowed; distal lobe spear-headed in shape, with a forceps-like outgrowth distally; lateral lobes slender and curved, minutely setose distally, shorter than the basal lobe.

Holotype, one male from Chengkiang, Yunnan Province, April 26, 1939, collected by the author. Allotype, one female from the same locality and date as the holotype. Paratype, one hundred and forty specimens with the same locality and date as the holotype; sixty specimens from the same locality, September, 1939, collected by the author; twenty specimens from the same locality, January, 1940, collected by the author. All types in the insect collection of Sun Yatsen University.

***Ochthebius (Bothochius) flexus, sp. nov.***

Size, 1.82 mm. long, 0.84 mm. wide.

Dorsal surface covered with blue procumbent hairs, head shining, dark brown, with metallic sheen; fronto-vertex punctured, intervals shining and generally wider than the punctures; a pair of large and deep depressions between the compound eyes, a small median depression behind; epistomal suture arched; clypeus punctured as fronto-vertex; labrum large, roughened, punctured and densely pubescent, more or less semicircular, anterior margin nearly straight; maxillary palpi brown, short, penultimate segment large and long, ultimate segment tapering distally; mentum brown, more or less trapezoid, sparsely and irregularly punctured.

Pronotum dark brown, punctured as fronto-vertex; median longitudinal furrow moderately deep with dilated anterior portion and tapering extremities; anterior and posterior discal depressions somewhat elliptical; outer depressions elongated, large and deep; anterior margin slightly emarginated, lined with a narrow translucent membrane; lateral margin retracted posteriorly from the anterior third, retracted portion lined with translucent membrane which encircles the posterior angle; posterior margin arched medially; anterior and posterior angles broadly rounded.

Scutellum black, triangular, glittering.

Thoracic sterna dull black, roughened and pubescent; prosternum slightly carinate longitudinally; mesosternum extending posteriorly between the middle coxae; metasternum large.

Elytra brown, moderately coarsely and deeply punctured, some punctures arranged in longitudinal direction; intervals shining, with minute lines and generally wider than the punctures; inner and outer margins with minute elevated border, those of the former more prominent posteriorly; humerales and lateral channels moderately prominent; anterior angles moderately sharp, posterior angles broadly rounded.

Legs brown; anterior coxae peg-like, pubescent and granulate; middle coxae nearly globular, pubescent and granulate; femora sparsely and setigerously punctured; tibiae shining, nearly straight, thickened distally, spinose; distal segment of tarsi much longer; claws slender and curved.

Five proximal abdominal sterna dull black, roughened and pubescent, narrow; sixth sternum dark brown, narrow, minutely punctured and pubescent posteriorly, proximal area roughened; seventh sternum dark brown, shining, punctured, posterior margin arched, minutely notched medially.

Basal lobe of phallus strongly curved, constricted at the middle, distal extremity pointed; distal lobe short, rounded distally; lateral lobes curved, distal ends with short setae.

Holotype, one male from Lou-fong-tsouen, Yunnan Province, March 26, 1940, collected by the author. Allotype, one female, with the same locality and date as the holotype. Paratype, seven specimens with the same locality and date as the holotype. All types in the insect collection of Sun Yatsen University.

***Ochthebius (Bothochius) lobatus*, sp. nov.**

This species is very close to *O. flexus*, but separable from it by the larger size (2.16 mm. long, 1.08 mm. wide), by the less prominent punctures on the head and the pronotum, by the deeper notch on anterior margin of labrum and by the absence of distinct minute lines on the interstices of elytra. The distal end of the basal lobe of the phallus of this species has the two sides equally convergent, while that of *O. flexus* has the two sides unequally convergent and the distal lobe of this species is in the form of a more or less circular lobe while that of *O. flexus* is in the form of an elongated lobe.

Holotype, one male from Chungking, Szechuan Province, August 8, 1942, collected by Miss S. C. Chen. Allotype, one female with the same locality and date as the holotype. Paratypes, twenty-six specimens with the same locality and date as the holotype. All types in the insect collection of Sun Yatsen University.

***Ochthebius (Hymenodes) exilis*, sp. nov.**

Size, 1.80 mm. long, 0.80 mm. wide.

Body color dark brown, with metallic shining, dorsal surface covered with moderately long procumbent hairs. Head dark brown, with large and sparse indistinct punctures; fronto-vertex and clypeus with minute dense granules; two large more or less elliptical deep depressions between the compound eyes and a smaller median one behind; epistomal suture slightly arched; clypeus somewhat trapezoid; labrum semicircular, densely and moderately coarsely punctured, with dense long hairs; maxillary palpi brown, penultimate segment large and long, ultimate segment slightly tapering distally; mentum dark brown, somewhat rectangular, anterior margin moderately deeply emarginated at the middle and slightly irregularly sculptured and punctured, intervals shining.

Pronotum dark brown, disc very minutely granulated and moderately sparsely and minutely punctured; median longitudinal furrow attenuated posteriorly, with tapering extremities; anterior discal depressions nearly rounded, posterior discal depressions elliptical; outer depressions large; anterior margin nearly straight; lateral margin retracted posteriorly from the point just in front of the middle, lined with translucent membrane which encircles the posterior angles; posterior margin arched; anterior angles rectangular, posterior angles rounded.

Scutellum dark brown, triangular, shining.

Thoracic sterna black, roughened; prosternum finely pubescent; mesosternum pubescent laterally, the median area raised into a long and triangular plate extending posteriorly between the middle coxae; metasternum densely pubescent, the posterior margin with a small notch medially.

Elytra dark brown; each elytron with ten series of punctures; interstices slightly raised, subequal to or slightly broader than the punctures in width and minutely granulated; inner margin with minute glittering border; lateral margin slightly arched, with minute elevated border; humerales prominent; anterior angles rectangular, posterior angles rounded.

Legs brown; anterior and middle coxae, more or less peg-like, roughened, pubescent; femora irregularly and setigerously punctured; tibiae thickened distally, strongly spinose; distal segment of tarsi long, subequal to the preceding segments together; claws slender and curved.

Five proximal abdominal sterna dull black, narrow, roughened and pubescent; sixth sternum broader, dark brown, shining, roughened; seventh sternum small; several slender setae at the distal end of the last abdominal tergum.

Basal lobe of the phallus long and curved; distal lobe short, blade-like; lateral

lobes slender, shorter than the basal lobe, distally with minute setae.

Holotype, one male from Lou-fong-tsouen, Yunnan Province, March 20, 1940, collected by the author. Allotype, one female with the same locality and date as the holotype. Paratype, eight specimens with the same locality and date as the holotype. All types in the insect collection of Sun Yatsen University.

***Ochthebius (Hymenodes) octofoveatus, sp. nov.***

Size, 1.68 mm. long, 0.80 mm. wide.

Head dark brown, distinctly granulated, with metallic shining; fronto-vertex with several shallow punctures; two large depressions between the compound eyes and a smaller median triangular depression behind, a small swelling present at the postero-lateral corner of each depression, median part of fronto-vertex slightly elevated as a rectangular area; epistomal suture arched; clypeus transverse; labrum indistinctly granulated, punctured, anterior margin with a semicircular deep notch at the middle; maxillary palpi dark brown, penultimate segment large, somewhat clavate, ultimate segment with tapering distal end; mentum brown, somewhat rectangular, roughened, minutely unevenly punctured, the anterior margin slightly emarginated.

Pronotum dark brown, disc evenly and minutely granulated; median longitudinal furrow with tapering extremities and separated into two parts by a median small swelling; anterior discal depression somewhat elliptical; posterior discal depression elongated, slightly oblique; two small discal depressions exterior to the anterior discal depressions and two larger ones exterior to the posterior discal depression; outer depressions large and long; anterior margin nearly straight, with a narrow translucent membrane, lateral margin retracted posteriorly from the anterior third, the retracted portion lined with membranous border; anterior angles somewhat rectangular, posterior angle broadly rounded.

Scutellum triangular, finely granulated.

Prosternum and mesosternum dark brown, granulated, the former carinated along the middle, extended posteriorly between the front coxae medially, the latter extended posteriorly between the middle coxae as a triangular plate; metasternum brown, roughened, very thinly pubescent, minutely notched posteriorly.

Elytra dark brown, each elytron with ten series of punctures; interstices glabrous, slightly raised and generally broader than or subequal to the punctures; inner margin with a minute glittering border; lateral margin arched, with minute elevated border; lateral channels and humerales prominent; anterior angles rounded, posterior angles broadly rounded.

Legs brown; front and middle coxae peg-like, roughened, thinly pubescent; femora and tibiae with minute sculpture; tibial spines moderately dense; distal segment of tarsi large and long; claws long and curved.

Abdomen brown, roughened; five proximal sterna narrow, thinly pubescent; sixth sternum broader, the median area shining and minutely irregularly punctured, roughened and thinly pubescent proximo-laterally, irregularly and minutely punctured and thinly pubescent disto-laterally; seventh sternum somewhat semicircular, shining, minutely and irregularly punctured, the proximal area with indistinct sculptured.

Holotype, one female from Lou-fong-tsouen, Yunnan Province, March 26, 1940, collected by the author, in the insect collection of Sun Yatsen University.